

Spin Dynamics in the Organic Quantum Magnet (C₅H₁₂N)₂CuBr₄

B. Thielemann¹, Ch. Rüegg², H. M. Rønnow¹, J. Stahn¹, J. Mesot¹,
D. McMorrow², K. W. Krämer³ and H.-U. Güdel³

(1) *Laboratory for Neutron Scattering, ETH Zürich & Paul Scherrer Institut,
5232 Villigen PSI*

(2) *Department of Physics and Astronomy and London Centre for
Nanotechnology, University College London, London WC1E 6BT*

(3) *Department of Chemistry and Biochemistry, University Bern,
3000 Bern 9*

The spin dynamics in the magnetic field versus temperature phase diagram of gapped quantum magnets, like Haldane-, ladder- and dimer-spin systems, has recently attracted considerable interest due to the presence of several novel quantum phases. Our search for a new model compound, where the complete phase diagram is accessible by neutron scattering experiments, has been successful. The potential organic ladder material (C₅H₁₂N)₂CuBr₄ has recently been grown in single crystalline form with sample masses (deuterated) suitable for such investigations. Its intrinsic energy scale with $H_{c1} = 6.6$ T and $H_{c2} = 13.5$ T, as determined from our inelastic neutron scattering experiments and magnetisation measurements, allows addressing for the first time the generic phase diagram of such a model spin system up to saturation. The system has therefore the highest potential for parametric studies. The spin dynamics in zero-field, including the determination of the spin Hamiltonian, its field-dependence, and the phase diagram for the region between H_{c1} and H_{c2} will be presented and compared to recent predictions from theory.